



**DR ADRIANA LE**  
ALLERGY | IMMUNOLOGY

**Dr Adriana Le**  
MBBS, FRACP  
Clinical Immunologist /Allergist

---

E [REDACTED]  
W allergyimmune.com.au

**Your Appointment**

Covid specialist vaccination

- Summit Health Centre, Mt Barker *clinic*
- Clare Medical Centre, Clare *18 707 49787*
- Health on Washington, Angaston



Government of South Australia  
SA Health

Walleroo Hospital  
Yorke and Northern Local  
Health Network

Ernest Terrace  
WALLAROO SA 5556

PMB 1  
WALLAROO SA 5556

Doctor Concerned  
Kadina Medical Associates  
PO Box 371  
Kadina SA 5554

Patient presented to Wallaroo Emergency Department. Please find a copy enclosed for your records

Administration  
Walleroo Hospital

# Dr Richard Daehn

FRACGP, B.Med.Sci: BMBS (Flinders)

Provider Number: [REDACTED]

Email: [REDACTED]

Argus: [REDACTED]

Healthlink: kadmedas

## Kadina Medical Associates

PO Box 371

Unit 77-79 Port Road

Kadina SA 5554

16 December 2021

Dr Adriana Le  
85 Wellington Road  
MOUNT BARKER 5251  
Ph: (08) 8406 7736  
Fax: (08) 8004 9151

Dear Dr Adriana,

Re:

Miss [REDACTED]

DOB:

Address:

Phone (Home):

Phone (Work):

Phone (Mobile):

Thank you for seeing Miss [REDACTED], age 50 yrs.

### Presenting Problem

In Association With:

Dr Graham Morris, Dr Michael Gregg, Dr Fazil Junejo, Dr Hannah Wellington, Dr Will Daehn

EFTA00263310

**Allergies**

Penicillins  
Lexapro  
Champix  
olanzapine

**Family History**

father high chol borderline diabetes

**Current Medications**

Nicotine 21mg per 24hour Transdermal Drug  
Delivery System (Transdermal patch)  
21mg/24hour(Step 1; 52.5 mg/patch) [28]

APPLY 1 NEW PATCH EVERY 24 HRS TO  
NONHAIRY, CLEAN, DRY SKIN ON  
TRUNK, UPPER ARM; HOLD IN PLACE  
FOR 10-20 SEC WITH PALM OF HAND;  
ROTATE APPLIC SITE (DO NOT  
REAPPLY TO SAME SITE WITHIN  
SEVERAL DAYS).

**PRN Medications**


Hiprex Tablets 1g [100]

1 tab twice daily

**Short Term Medications**

Should any of the above need clarification, please feel free to contact me by phone on 8821 3133, or e-mail me at [REDACTED]. Can you please inform me if this patient does not make an appointment or does not keep an appointment.

Yours sincerely,

  
.....  
**Dr Richard Daehn**



**DR ADRIANA LE**  
ALLERGY | IMMUNOLOGY

Immunologist/Allergist  
MBBS FRACP

Practices

Summit Health Centre  
85 Wellington Rd, Mt Barker 421536DF

Clare Medical Centre  
41 Old N Rd, Clare  
421536ET

Health on Washington  
3-7 Fife St, Angaston  
421536GL


**CONTACT**



W allergyimmune.com.au

04/02/2022  
Dr Richard Daehn  
Kadina Medical Associates  
PO Box 371  
KADINA SA 5554


Dear Richard,



Kind Regards,

Dr Adriana Le

This vaccine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. You can report side effects to your doctor, or directly at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).



# COMIRNATY™ COVID-19 VACCINE

Active ingredient: BNT162b2 [mRNA]

This vaccine has provisional approval in Australia to prevent COVID-19 disease caused by SARS-CoV-2 virus in adults and adolescents from 16 years of age and older. This approval has been granted on the basis of short term safety and efficacy data. Evidence of longer term efficacy and safety from ongoing clinical trials and vaccination in the community continues to be gathered and assessed.

## Consumer Medicine Information (CMI)

This leaflet provides important information about using COMIRNATY. You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about receiving COMIRNATY.

Where to find information in this leaflet:

1. [Why am I being given COMIRNATY?](#)
2. [What should I know before I am given COMIRNATY?](#)
3. [What if I am taking other medicines?](#)
4. [How will I be given COMIRNATY?](#)
5. [What should I know while being given COMIRNATY?](#)
6. [Are there any side effects?](#)
7. [Product details](#)

### 1. Why am I being given COMIRNATY?

COMIRNATY contains the active ingredient BNT162b2 [mRNA]. COMIRNATY is an mRNA (messenger ribonucleic acid) vaccine.

COMIRNATY is a vaccine given to prevent COVID-19 disease caused by SARS-CoV-2 virus in adults and adolescents from 16 years of age and older.

COMIRNATY works by triggering your immune system to produce antibodies and blood cells that work against the virus, to protect against COVID-19 disease.

### 2. What should I know before I am given COMIRNATY?

Warnings



COMIRNATY should not be given:

1. If you are allergic to BNT162b2 [mRNA] or any of the ingredients listed at the end of this leaflet.

Check with your doctor if you have:

- 
- had a severe allergic reaction or breathing problems after any other vaccine or after being given COMIRNATY in the past.
  - fainted following any needle injection.

- a severe illness or infection with high fever. However, you can have your vaccination if you have a mild fever or upper airway infection like a cold.
- a weakened immune system, such as due to HIV infection or are on a medicine that affects your immune system.
- a bleeding disorder, bruise easily or are on a blood thinning medicine.

As with any vaccine, COMIRNATY may not fully protect all those who receive it and it is not known how long you will be protected.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section 6. [Are there any side effects?](#)

Pregnancy and breastfeeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before you receive this vaccine.

Children and adolescents

COMIRNATY should not be given to children under 16 years.

### 3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Tell your doctor or pharmacist if you have recently received any other vaccine.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect, or are affected by, COMIRNATY.

### 4. How will I be given COMIRNATY?

- COMIRNATY will be given as an injection into the muscle of your upper arm by a doctor, nurse or pharmacist.

- You will be given one dose followed by a second dose at least 21 days later. If you miss the second dose, ask your doctor for advice.
- A doctor, nurse or pharmacist will observe you for at least 15 minutes after being given COMIRNATY.

## 5. What should I know while being given COMIRNATY?

### Things you should know

- If you receive one dose of COMIRNATY, you should receive a second dose of the same vaccine at least 21 days later to complete the vaccination schedule.
- You may not be protected against COVID-19 disease until at least seven days after your second dose.
- You may not be protected if you only receive one dose, so a second dose is important.

### Driving or using machines

Be careful before you drive or use any machines or tools until you know how COMIRNATY affects you.

Some of the side effects of COMIRNATY may temporarily affect your ability to drive or use machines.

### Storage of the vaccine

COMIRNATY is stored at -90 °C to -60°C. It must be kept in the original package in order to protect from light.

A doctor, nurse or pharmacist will prepare the injection for you before you are given it.

### Getting rid of any unwanted vaccine

A doctor, nurse or pharmacist will dispose of any unused vaccine.

## 6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

### Very common side effects

Very common side effects	What to do
<ul style="list-style-type: none"> <li>• pain/swelling at injection site</li> <li>• tiredness</li> <li>• headache</li> <li>• muscle pain</li> <li>• chills, fever</li> <li>• joint pain</li> </ul>	Speak to your doctor if you have any of these very common side effects and they worry you.

### Common side effects

Common side effects	What to do
<ul style="list-style-type: none"> <li>• redness at injection site</li> <li>• nausea</li> </ul>	Speak to your doctor if you have any of these common side effects and they worry you.

### Uncommon side effects

Uncommon side effects	What to do
<ul style="list-style-type: none"> <li>• enlarged lymph nodes</li> <li>• feeling unwell</li> <li>• pain in limb</li> <li>• insomnia</li> <li>• itching at injection site</li> </ul>	Speak to your doctor if you have any of these side effects and they worry you.

### Rare side effects

Rare side effects	What to do
<ul style="list-style-type: none"> <li>• temporary one-sided facial drooping</li> </ul>	Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice this side effect.

### Other side effects (frequency unknown)

Other side effects (frequency unknown)	What to do
<ul style="list-style-type: none"> <li>• severe allergic reaction</li> </ul>	Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice this serious side effect.

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

### Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side