

December 21, 2015

Anat Maoz Identity Card No. [REDACTED]

Age: 49, M + 4, Painter

Ethnicity: Ashkenazi

Family history: Both grandmothers had breast cancer at age 60+

Genetics: Negative for common mutations

Concomitant diseases: No

Regular medications: No

Gynecology history: Leukeran since 2013

**History of Present Illness:**

In 2006, at age 40, was diagnosed with advanced breast cancer without lymph node involvement. RS-23. Underwent lumpectomy and sentinel node resection. Received radiotherapy and Tamoxifen. Finished taking Tamoxifen in 2009. Since 2010 has been followed with serial imaging for suspected T1 metastases. In 2013 had a definitive diagnosis of T1 metastasis. Underwent surgery – fixation, focused radiation and resumed Tamoxifen therapy.

For more than a year she has been complaining that the right axilla does not feel right – no imaging studies.

In September, 2015, she developed right hand numbness. Underwent a systemic work-up which revealed:

1. Tumor spread to the right axillary lymph nodes (PET/CT and directed US)
2. T1 destruction with invasion into the canal (PET/CT and MRI)
3. Rise in tumor markers
4. No evidence of additional spread

Has begun taking Leukeran + Letrozole. Is scheduled to begin taking Xgeva.

There may be a slight improvement in the neurological complaints. Axillary discomfort has improved.

T1 surgery had been recommended followed by stereotactic radiation – at this point in time the patient is refusing surgery.

**On examination:**

In good state of health. I could not palpate with certainty any axillary lymph nodes.